

Contents lists available at [SciVerse ScienceDirect](http://www.sciencedirect.com)

Biochimica et Biophysica Acta

journal homepage: www.elsevier.com/locate/bbabio

S18 Supercomplexes / Proteomics

18L1

Identification of new assembly factors of complex I by complexome profiling

I. Wittig, H. Heide, L. Bleier, B. Schwamb, V. Strecker, M. Steger, J. Ackermann, S. Dröse, M. Zörnig, A.S. Reichert, I. Koch, U. Brandt
 Molecular Bioenergetics Group, Medical School, Cluster of Excellence Frankfurt "Macromolecular Complexes," Goethe-University, Theodor-Stern-Kai 7, D-60590 Frankfurt am Main, Germany
 Chemotherapeutisches Forschungsinstitut Georg-Speyer-Haus, Paul-Ehrlich-Straße 42-44, D-60596 Frankfurt am Main, Germany
 Molecular Bioinformatics Group, Institute of Computer Science, Faculty of Computer Science and Mathematics, Robert-Mayer-Strasse 11-15, D-60325 Frankfurt am Main, Germany
 Mitochondrial Biology, Buchmann Institute for Molecular Life Sciences, Goethe University Frankfurt am Main, Max-von-Laue-Str. 15, D-60438 Frankfurt am Main, Germany
 Mitochondriale Biologie, Zentrum für Molekulare Medizin, Goethe Universität Frankfurt am Main, Max-von-Laue-Str. 15, D-60438 Frankfurt am Main, Germany
 E-mail: wittig@zbc.kgu.de

Blue-native electrophoresis (BNE) is commonly used for the isolation and quantification of OXPHOS complexes. In order to study composition and dynamics of lower abundant mitochondrial complexes we combined BNE and label free quantitative mass spectrometry to analyze the complexome of isolated rat heart mitochondria. Hierarchical clustering of proteins with similar migration in BN-lanes allowed the identification of well-characterized mitochondrial complexes and unknown protein–protein interactions. The known complex I assembly factors CIA30, ECSIT and ACAD9 were found in a cluster together with TMEM126B, a protein of unknown function. Stable lentiviral knockdown of TMEM126B in the osteosarcoma cell line 143B resulted in a strong reduction of complex I levels without affecting the assembly of other OXPHOS complexes. Functional characterization by high-resolution respirometry of intact mitochondria from knockdown cells revealed a reduction of the complex I dependent respiration by roughly 65%. In contrast, succinate dependent respiration was unaffected. In addition, we characterized the homologous protein TMEM126A that was recently reported to be associated to mitochondrial diseases. Knockdown cells showed complex I deficiency but TMEM126A was not identified within the mitochondrial complex I assembly factor (MCIA) complex containing CIA30, ACAD9, ECSIT and TMEM126B. In summary, complexome profiling is a powerful bottom-up approach allowing identification of unknown protein interactions of an entire organelle.

doi:[10.1016/j.bbabio.2012.06.363](https://doi.org/10.1016/j.bbabio.2012.06.363)

18P1

Study of respiratory complexes in the low GC firmicutes *Bacillus subtilis* and *Geobacillus stearothermophilus*

L. Bergdoll, E. Point, F. Bayman, D. Picot
 Laboratoire de Physico-Chimie Moléculaire des Membranes Biologiques UMR7099 CNRS-Université Paris Diderot, Institut de Biologie Physico-Chimique, 13 rue Pierre et Marie Curie, 75005 Paris, France
 BIP, Centre National de la Recherche Scientifique, UMR7281, Centre National de la Recherche Scientifique/Aix Marseille University, FR3479, 31 Chemin Joseph Aiguier, Marseille, France
 E-mail: lucie.bergdoll@ibpc.fr

Rieske/cytochrome *b* complexes play a central role in energy production by translocating electrons and protons across the membrane, building up an electrochemical potential. This enzyme family is composed of several members, among them cytochrome *bc*₁ and *b*₆*f* complexes which have been shown to distinguish themselves with respect to the structure of their c-type cytochrome subunit and the co-factor content. Importantly an additional heme, heme *c*_i, is present in the quinone reduction side of *b*₆*f* complexes. Its function is not yet understood.

Here, we set out to study protein/protein and protein/lipid interfaces on the Rieske/cytochrome *b* complex from *Geobacillus stearothermophilus*, a thermophilic low GC gram-positive bacterium. This enzyme is integrated in the respiratory chain of the bacterium and more closely related to cytochrome *b*₆*f* from photosynthetic reaction chains than to cytochrome *bc*₁ from the mitochondrial respiratory chain. At variance with mitochondrial *bc*₁ and chloroplasts *b*₆*f* complexes the enzyme of *G. stearothermophilus* uses menaquinone as a substrate. This complex has been enriched by anion exchange chromatography and density gradient centrifugation. Here we present a first characterization by UV/VIS and EPR spectroscopy.

In parallel, we started studying enzymes of the respiratory chain from *Bacillus subtilis*, a mesophilic bacterium, which is closely related to *G. stearothermophilus*. The respiratory complexes of these two organisms are quite similar, but exhibit different molecular organizations and stabilities.

References

- [1] W. Nitschke, R. Van Lis, B. Schoepp-Cothenet, F. Baymann, The "green" phylogenetic clade of Rieske/cytb complexes, Photosynth. Res. (2010).
- [2] D. Stroebel, Y. Choquet, J.L. Popot, D. Picot, An atypical heme in the cytochrome *b*₆*f* complex, Nature 426 (2003) 413–418.

doi:[10.1016/j.bbabio.2012.06.364](https://doi.org/10.1016/j.bbabio.2012.06.364)